

International Concurrent Workshop Clinical 1

RA biomarker

Day 1 April 21, 9:10 - 10:10 Room 4

Chair: Toshihiro Nanki

Division of Rheumatology, Department of Internal Medicine, Toho University School of Medicine, Japan

ICW-C1-1 HLA-DRB1 analysis utilizing consecutive data of rheumatoid factor (RF) identified a genetically unique subset among rheumatoid arthritis and distinct genetic background of RF levels from anti-cyclic citrullinated peptide antibodies

Ryosuke Hiwa

Department of Rheumatology and Clinical Immunology, Graduate School of Medicine, Kyoto University, Japan

ICW-C1-2 CIGARETTE SMOKE TRIGGERED CIRCULATING AND INFILTRATING NEUTROPHIL EXTRACELLULAR TRAPS BY AUTOPHAGY DEPENDENT WAY IN RA PATIENTS

Li Wang

Rheumatology and Immunology, Affiliated Anhui Provincial Hospital, Anhui Medical University, Hefei, Anhui, China

ICW-C1-3 14-3-3 η is associated with immunological and structural status as well as disease activity in rheumatoid arthritis

Shintaro Hirata

The First Department of Internal Medicine, School of Medicine, University of Occupational and Environmental Health, Japan / Department of Laboratory and Transfusion Medicine, University Hospital, University of Occupational and Environmental Health, Japan

ICW-C1-4 Predictive value of the multi-biomarker disease activity (MBDA) score for flare and sustained remission in the HONOR study

Shintaro Hirata

The First Department of Internal Medicine, School of Medicine, University of Occupational and Environmental Health, Japan / Department of Laboratory and Transfusion Medicine, University Hospital, University of Occupational and Environmental Health, Japan

ICW-C1-5 What blood inflammatory marker should be the most influential on patient discomfort among rheumatoid arthritis patients treated with Infliximab or Tocilizumab?

Kazuha Kizaki

Department of Orthopaedic Surgery and Rheumatology, Kyoto Shimogamo Hospital, Japan

ICW-C1-6 Presepsin (soluble CD14 subtype) and procalcitonin as biomarkers of systemic bacterial infection in patients with rheumatoid arthritis

Kohei Tsujimoto

Department of Clinical Immunology and Rheumatology, Kitano Hospital, The Tazuke Kofukai Medical Research Institute, Japan