

International Concurrent Workshop Clinical 10

Myopathy

Day 3 April 23, 8:30 - 9:30 Room 4

Chair: Hideto Kameda

Division of Rheumatology, Department of Internal Medicine, Toho University Ohashi Medical Center, Japan

ICW-C10-1 The availability of Nailfold Videocapillaroscopy (NVC) in patients with polymyositis and dermatomyositis (PM/DM)

Yasuyuki Todoroki

The 1st Department of Internal Medicine, University of Occupational and Environmental Health, Japan

ICW-C10-2 Inflammatory myositis associated with malignancy: A retrospective multicenter study of 213 Japanese patients with Dermatomyositis and Polymyositis

Shinichiro Omura

Nagoya City University Hospital, Japan

ICW-C10-3 Clinical course and treatment of two refractory cases of anti-HMGCR antibody-associated necrotizing autoimmune myopathy

Sumie Hiramatsu

Department of Nephrology, Rheumatology, Endocrinology and Metabolism, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Japan

ICW-C10-4 Assessment of peripheral joints by musculoskeletal ultrasonography in polymyalgia rheumatica patients predicts future DMARDs use

Naomi Tsuchida

Department of Internal Medicine and Clinical Immunology, Yokohama City University Graduate School of Medicine, Yokohama, Japan

ICW-C10-5 Tocilizumab (TCZ) but not methotrexate (MTX) is steroid-sparing in polymyalgia rheumatica (PMR): A single-center retrospective study

Keisuke Izumi

Department of Connective Tissue Diseases, National Tokyo Medical Center, Japan / Division of Rheumatology, Department of Internal Medicine, Keio University School of Medicine, Japan

ICW-C10-6 The glucocorticoid-sparing effect of tocilizumab (TCZ) is superior to that of methotrexate (MTX) in polymyalgia rheumatica (PMR): A singlecenter retrospective study

Okinori Murata

Division of Rheumatology, Department of Internal Medicine, Keio University School of Medicine, Tokyo, Japan